AD	

Award Number: DAMD17-00-1-0063

TITLE: A Re-examination of Neuropsychological Functioning in

Persian Gulf War Era Veterans

PRINCIPAL INVESTIGATOR: Roberta F. White, Ph.D.

CONTRACTING ORGANIZATION: Boston University

Boston, Massachusetts 02118

REPORT DATE: February 2001

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;

Distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED		
	February 2001	Annual (24 Jan	00 - 23 Ja	an 01)
4. TITLE AND SUBTITLE			5. FUNDING N	UMBERS
A Re-examination of	Neuropsychological	L	DAMD17-00-	-1-0063
Functioning in Persian Gulf War Era Veterans				
			-	
6. AUTHOR(S)				
Roberta F. White, Ph.D.				
				A A SAN AND A SA
7. PERFORMING ORGANIZATION NAM	ME(S) AND ADDRESS(ES)		8. PERFORMING REPORT NUI	G ORGANIZATION
Boston University			REPORT NO	WIDER
Boston, Massachusetts 02118				
E-MAIL: rwhite@bu.edu	TION NAME (C) AND ADDRESS (FC	1	10 CDONCODII	NC / MONITORING
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
MG A M Park Dance to and A	fatarial Commond			
U.S. Army Medical Research and Materiel Command				
Fort Detrick, Maryland 21702-501	2			
11. SUPPLEMENTARY NOTES				
THE GOTT ELINERT ATT. NOTES				
12a. DISTRIBUTION / AVAILABILITY S				12b. DISTRIBUTION CODE
Approved for public rele	ease; Distribution unl	imited		
1				

13. ABSTRACT (Maximum 200 Words)

Existing data from our group suggest that Gulf War (GW) veterans score worse on a specific group of neuropsychological tests than GW-era veterans not deployed to the Gulf. These findings raise the possibility of subtle, "subclinical" central nervous system (CNS) damage associated with Gulf deployment. A critical question is whether these findings may be attributable to exposure to neurotoxicants in the Gulf environment.

Recently some PGW veterans have reported that their cognitive functioning has been <u>worsening</u> over the time since their return from the Gulf. This report is at odds with the usual course of CNS effects of intoxication, which generally remit or remain static in the absence of exposure. The reports of declining cognition raise the issue of exposure to toxicants that might produce a progressive condition versus the existence of other conditions that might affect cognition in individuals who are experiencing progressive functional declines.

The specific aims of the project are: 1) to determine whether objective test measures reveal progressive diminution in cognitive function by comparing test performance observed initially (1995-1998) to performance 4 years (+/- 6 months) later and 2) to identify the factors associated with any cognitive diminutions including chemical exposures, stress, or psychiatric disorders.

14. SUBJECT TERMS Gulf War Illness	15. NUMBER OF PAGES -42		
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. Z39-18 298-102

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

PI - Signature

Date

TABLE OF CONTENTS

Front Cover	1
SF 298	2
Foreword	3
Table of Contents	4
Introduction	5
Body of report	5
Preliminary Studies	14
Key Research Accomplishments	30
Reportable Outcomes	
References	34
Appendixes	4

INTRODUCTION

Our research completed to date indicates that Persian Gulf War (PGW) veterans deployed to the Gulf score more poorly on a limited, specific group of neuropsychological test measures than PGW-era veterans who were not deployed to the Gulf. These findings raise the possibility of subtle, "subclinical" central nervous system damage associated with Gulf deployment. The etiology of the neuropsychological deficits is unclear, but our data suggest that they are related to exposures experienced in the Gulf. Our data also suggest that these deficits cannot be attributed to stress, psychiatric status, or compensation seeking.

Recently some PGW veterans who were deployed to the Gulf have reported clinically that their cognitive functioning has been worsening over the time since their return from the Gulf. Specifically, study participants report declines in the areas of short-term memory and concentration. Additionally, research subjects with high symptom complaints continue to report ongoing high symptom levels (SPProctor, personal communication). These reports are somewhat at odds with the <u>usual course</u> of central nervous system (CNS) effects of intoxication, which generally remit or remain static in the absence of exposure.

The specific aims of this project are: 1) to determine whether objective test measures reveal any progressive diminution in cognitive function by comparing test performance observed initially (1995-1998) to performance 4 – 5 years later among Gulf-deployed veterans compared to non-Gulf-deployed controls, and 2) to determine whether any identifiable declines in function are related to Gulf-related environmental exposures, stress, psychiatric disorders, existence of multiple health complaints, normal aging, or motivational disorders.

Body

Background and Significance.

Environmental Exposures and Health Effects

The most commonly reported health related illnesses of Persian Gulf War (PGW) veterans included fatigue, headache, skin rash, joint or muscle pain, and loss of memory or other cognitive difficulties (Institute of Medicine report, 1996). The Department of Defense Comprehensive Clinical Examination Program (CCEP) data set further identified these symptoms as well as sleep disturbance, concentration problems, cough, and

depression as commonly reported by a large study sample of 18,000 respondents (Roy, Chung 1994; Institute of Medicine report 1996). In studies conducted through the Boston Environmental Hazards Center it has been found that veterans who were deployed to the Gulf reported the most prominent symptoms to be forgetfulness, fatigue, restless sleep, gastrointestinal disturbance, and pulmonary and cardiac symptoms (Proctor et al., 1998). In 1994, medical examinations of active duty PGW veterans indicated that the most common diagnoses of these veterans included diseases of the musculoskeletal system and connective tissue, mental disorders, ill-defined conditions, skin diseases, digestive system diseases, diseases of nervous system/sensory organs and infectious and parasitic diseases (CCEP, Joseph, 1997). A study of hospital discharges in 1992 and 1993, indicated no significant difference in overall risk for hospitalization in PGW veterans compared with non-PGW veterans. However, the specific diagnoses that were hospitalized more frequently in PGW veterans included neoplasms, genitourinary diseases, blood diseases, and mental disorders (Gray et al., 1996). Longitudinal surveys of a large cohort of GW veterans have suggested that many veterans are reporting their health symptoms to be gradually worsening since their original evaluations shortly after their return from the Gulf War theater (SP Proctor, personal communication).

Persian Gulf War veterans may have been exposed to many different environmental toxicants. Some researchers suggest that this may be the cause of their reported health complaints (US Senate Committee Rockefeller Report, 1995; Haley, 1997). These possible exposures include smoke from burning Persian Gulf oil wells (US Congress, Office of Technology Assessment, September, 1993), biological or chemical nerve agents deployed by Iraq (Riegle, 1993), pre-deployment vaccines, anti-nerve gas agents (e.g., pyridostigmine), skin absorption of diesel fuel, airborne or material-based organic solvents, pesticides, parasitic infection (e.g., leishmaniasis tropica) and radiation from depleted uranium weaponry. Sarin, a known neurotoxicant, (Kadar et al., 1995) and mustard gas were reportedly detected in Saudi Arabia (Washington Post, Boston Globe, 11/1/93 and Boston Globe, 11/24/93). In addition, the US Department of Defense identified a chemical weapons arsenal at a destroyed storage site at Khamisiyah, Iraq. US military personnel charged with overseeing the destruction were unaware of the

chemical weapons in this site and appropriate precautions were not taken against such exposures. In January 2001, the Pentagon revised its numbers of potentially exposed veterans by an additional 35,000 individuals. At other sites, measurements by US chemical warfare vehicles also reported unconfirmed measurement of the mustard gas lewisite as well as an unidentified blistering agent and benzyl bromide.

Sequelae from exposure to these environmental toxicants often include damage to multiple organ systems; for example, lewisite and mustard gases produce damage to the skin, respiratory system, and appear to be potent carcinogens (e.g., Chevillard, Lainee, Robineau, & Puchelle, 1992; Pechura & Rall, 1993; Watson, Jones, & Griffin, 1989). Polycyclic aromatic hydrocarbons (PAHs) from oil smoke generate in vitro genotoxicity, adversely impacting immune and respiratory functions (e.g., Klaassen, Amdur, & Doull, 1986; Kelsey et al., 1994) and depleted uranium has been cited as having carcinogenic effects in addition to adverse effects on kidney functioning (Doucet, 1994). Other purported environmental exposures such as kerosene and diesel fuel (Andrews & Snyder, 1986); pyridostigmine (Almog et al., 1991; Keeler, Hurst, & Dunn, 1991); metals and volatile organics found in smoke (Klaassen et al., 1986); and sarin (Burchfiel & Duffy, 1982; Kawabuchi, Cintra, Deshpande, & Dlbuquerque, 1991; Husain, Vijayaraghavan, Pant, Raza, & Pandey, 1993). exert both acute and chronic neurotoxicant effects in animals and humans. Although, an incidence study of admissions to an Army hospital during the PGW reported no admissions for exposure to chemical or biological weapons (Wintermeyer, 1994), it is possible that chronic low level exposures may be the cause of cognitive and physiological changes as long-term sequelae are unknown in humans. In fact, a recent report suggested that low-level neurotoxicant exposures may cause the delayed-onset neurotoxic syndrome OPIDP (organophosphate-induced delayed polyneuropathy) that may account for the varied symptoms of GW veterans (Kurt, 1998).

Correspondingly, neuropsychological evaluations are the instrument of choice for low-level neurotoxicant exposures because these tests can not only quantify behavioral deficits in persons with clinically obvious physical evidence of intoxication following exposure (Baker & White, 1985) but can also detect subtle dysfunction in exposed persons without obvious clinical disease. That is, even in the absence of overt toxicant

poisoning, neurotoxicant exposed workers have been found to score significantly worse on specific neuropsychological tests than non-exposed workers. This condition has been referred to as "subclinical encephalopathy" (Baker, White & Murawski, 1985; White & Feldman, 1987).

Particularly important with regard to the possibility of identifying the source of toxicant exposure in a situation with multiple possible agents (such as in the PGW) is the specificity of relations between particular toxicants and distinct neurobehavioral and neuropsychological effects (White et al., 1992). Epidemiological studies during the past 30 years have examined the impact of exposure to metals (e.g., lead, mercury, arsenic), organic solvents (e.g., trichloroethylene, n-hexane, petroleum distillates), and pesticides (e.g., organophosphates, carbamates) on cognitive functioning. For example, studies of solvent exposure have reliably shown disturbances in executive function, attention, visuospatial skills, short-term memory, and mood (e.g., Anger, 1990; Echeverria & White, 1992; White, Feldman & Proctor, 1992). Studies of lead-exposed workers have yielded similar findings along with decrements in verbal reasoning and motor functions (e.g., Baker et al., 1984; Hanninen, Hernberg, Mantere, Vesanto, & Jalkaner, 1978; Yokoyama, Araki, & Aono, 1988).

Neuropsychological functioning in Persian Gulf War Veterans

Recent neuropsychological comparisons of PGW veterans have shown significant differences when compared with controls or when compared with published normative data (Goldstein 1996; Axelrod et al, 1997, Homm, 1997). These studies showed significant differences in PGW veterans on measures of executive system functioning (Stroop test), two measures of motor functioning (Finger tap test, Purdue Pegboard test), and several overall impairment indexes including items from the Pittsburgh Occupational Exposure Battery and WAIS-R full scale IQ scores. It should be noted however, that some of these studies included small group sizes (Goldstein, 1996, Hom et al., 1997) and did not include appropriate comparison groups (Axelrod, 1997). An additional study suggested that emotional functioning was correlated with observed cognitive deficits in PGW veterans such that sustained

attention was associated with trait anxiety, and depression was associated with motor coordination and executive system functioning (Sillanpaa et al. 1997). Hom et al., (1997) suggested from their findings that sick GW veterans showed generalized cognitive deficits rather than specific cognitive domain deficits. This pattern of cognitive deficit is not the usual course expected from mild neurotoxicant exposures (White & Proctor, 1997) and if replicated raises questions about multiple neurotoxicant exposures in these veterans.

Existing data from our group (White et al., in press) suggest that Gulf War veterans score more poorly on a limited, specific group of neuropsychological test measures than PGW-era veterans who were not deployed to the Gulf. These findings raise the possibility of subtle, "subclinical" central nervous system damage associated with Gulf deployment. A critical question about these findings is whether they may be attributable to exposure to neurotoxicants in the Gulf environment. This possibility is particularly salient because the existence of cognitive deficits was related in our sample to self-reported exposures to substances in the Gulf theater that are known to be neurotoxic.

Recently some PGW veterans have reported that their cognitive functioning has been worsening over the time since their return from the Gulf. This report is at odds with the usual course of central nervous system (CNS) effects of intoxication, which generally remit or remain static in the absence of exposure. The reports of declining cognition raise the issue of exposure to toxicants that might produce a progressive condition versus the existence of other conditions that might affect cognition in individuals who are experiencing progressive functional declines.

Effects of Stress on Physical Health

The relationship between physical health symptoms and psychological stressors in PGW veterans has raised several important questions regarding illness mechanisms. For instance, it has been found that stress and subsequent physiological arousal and anxiety is associated with subsequent reports of deteriorating health status, including cognitive declines in memory, concentration, and attention (e.g.,

Shalev, Bleich, & Ursano, 1990; Wolfe et al., 1992). Other researchers suggest that prolonged stress can lead to long-term hormonal changes (Yehuda, 1997), reduced immune system functioning, (Black, 1994) and alterations of brain neurotransmitters that mediate cognitive and psychological responses (Wolfe & Charney, 1991). Reports of animal models suggest that acute stress may also alter the permeability of the blood brain barrier to particular neurotoxicants (including pyridostigmine bromide; an acetylcholinesterase inhibitor given prophylactically to PGW veterans as a protective measure in the event of chemical weapon attack (Friedman et al 1996). However, other reports of different rodent species could not replicate these findings (Lallement, 1998). Findings from our original evaluations of treatment seeking GW veterans suggested that self-reported pyridostigmine bromide (PB) use was associated with lower scores on attention/executive system functioning (Wisconsin card sort test) and verbal memory functioning (California Verbal Learning Test) compared with veterans not reporting PB use. However, we were not able to confirm an interaction effect of reported PB use and severe stress reactions such as PTSD diagnosis (Sullivan et al., 2001).

To date, the symptoms reported by PGW veterans have received no definitive diagnosis, and may not constitute a single diagnostic entity. In fact, a factor analysis by Haley et al. (1997) identified 6 possible syndrome types in this population that may represent delayed-onset neurotoxic syndrome variants. These subtypes include impaired cognition, confusion-ataxia, arthro-myo-neuropathy, phobia-apraxia, fever-adenopathy and weakness-incontinence. Others however, after noting symptom complaints of fatigue, mood and musculoskeletal complaints in PGW veterans, suggest that half of these veterans would satisfy the criteria for chronic fatigue syndrome (Reeves, 1995). However, our original evaluation with 200 GW veterans found <1 percent met clinical criteria for chronic fatigue syndrome. In addition, other researchers suggest that infectious diseases are the cause of the varied symptoms noted by these veterans (Nicolson et al, 1996; Oster et al., 1992; Kreutzer et al, 1993). Drug treatments are currently underway to assess if large doses of antibiotics may reduce complaints in veterans testing positive for mycoplasmic infections. Cross-sectional studies of veterans suffering from post-traumatic

stress disorder (PTSD) have found that individuals with this diagnosis consistently report more physical health symptoms and poorer health status overall than their non-PTSD counterparts (Kulka et al., 1990; Litz et al., 1992; Shalev et al., 1990; Kimerling & Calhoun, 1994; Wolfe et al., 1993). In addition, these veterans also showed lower salivary cortisol levels than their non-PTSD counterparts suggesting dysregulated immune system functioning (Kellner et al., 1999). However, we were unable to confirm any health complications from the synergistic effects of small doses of neurotoxicants combined with a stress induced reduction in immune functioning (Friedman et al, 1996) in our original evaluation of 200 GW veterans. In sum, there are several competing hypotheses for the varied symptoms reported by PGW veterans. These etiologies include environmental exposures, post-traumatic stress disorder, chronic fatigue syndrome and multiple chemical sensitivity.

Post-traumatic Stress Disorder

The essential feature of Post -Traumatic Stress Disorder (PTSD) is the development of characteristic symptoms following a psychologically traumatic event that is generally outside the range of usual human experience. The characteristic symptoms involve re-experiencing the traumatic event; numbing of responsiveness to or reduced involvement with the external world; and a variety of autonomic, dysphoric, or cognitive symptoms (American Psychiatric Association, 1995).

Rates of presumptive PTSD in PGW veterans vary considerably across samples (3-46%), dependent in part upon unit mission, unit composition, and location in the Middle East according to data from a longitudinal study of nearly 3,000 PGW veterans (Wolfe et al., 1992; Wolfe et al., 1993). A recent meta-analysis by Haley (1997) suggested that incidence rates of PTSD in PGW veterans were overestimated in some studies. Cognitive performance has been noted to be compromised in subjects with PTSD and include problems with attention/mental tracking (e.g., span of apprehension) and executive system functioning (e.g., planning, anticipatory skills, monitoring), but not to problems with

acquisition of information, retroactive interference, mental manipulation and sustained attention in PGW veterans diagnosed with PTSD. Depression (rather than PTSD) is associated only with attention and mental tracking problems. By comparison, elements of physical trauma in prisoners of war (e.g., percent captivity weight loss) show a significant association with learning and memory deficits. Overall, increased severity of PTSD, depression, and weight loss are associated with greater rates of cognitive dysfunction (see also Levy, 1988). Our initial evaluations suggested no significant cognitive differences in performance of GW veterans with PTSD compared with GW veterans without a PTSD diagnosis. However, veterans with PTSD reported significantly more mood disturbances than their non-PTSD counterparts and suggest the need to control for mood disturbances when conducting cognitive evaluations of veterans with PTSD.

Multiple Chemical Sensitivity and Chronic Fatigue Syndrome

Many of the symptoms reported by PGW veterans overlap with those associated with two other diagnoses whose etiology has not yet been determined: multiple chemical sensitivity (MCS) and chronic fatigue syndrome (CFS). A definition of MCS offered by Cullen (1987) has gained acceptance: symptoms triggered by an identifiable environmental exposure; present in more than one organ system; not explained on the basis of any standard test of organ system function; recurring and abating in response to exposure to chemicals of diverse structural classes that involve differing modes of action, at levels of exposure far below those typically known to produce adverse reactions in humans.

A definition of CFS offered by Fukuda et al. (1994) has gained acceptance: extreme fatigue lasting at least six months; producing a significant reduction in daily activities; not resolving with bed rest. The disorder may be accompanied by, but not predated by: self-reported short-term memory loss; sore throat; tender cervical or axillary lymph nodes; muscle pain; multi-joint pain without joint swelling/redness; headaches of a new type, pattern or severity; unrefreshing sleep; post-exertion malaise lasting more than 24 hours. The diagnosis requires four or more of the above symptoms. These

symptoms are commonly seen in PGW veterans. Exclusionary criteria include: an active medical condition, such as untreated hypothyroidism; obesity; a previously diagnosed medical condition whose resolution has not been documented; past or current diagnosis of major depression, bipolar affective disorder, dementia, or schizophrenia; alcohol or substance abuse within 2 years. Results of our original study suggested that although many GW veterans report MCS and CFS like symptoms, very few meet actual criteria for these diagnoses (<1 percent). Thus, it seems unlikely that GW syndrome can be attributed to either of these two diagnostic entities although all three illnesses may share some heretofore-unknown origin.

Overall Project Objective.

This study will focus primarily on longitudinal analyses and comparison of neuropsychological functioning among our previously examined GW-era cohort. We will, however, also examine differences in health symptom report, and diagnostic outcomes (e.g., post-traumatic stress disorder [PTSD], and psychiatric disorders). Neuropsychological functioning was chosen as a primary focus because of its relevance to the multiple CNS complaints reported by PGW veterans and because of the sensitivity of neuropsychological measures to the diagnoses of interest in this situation. We will specifically explore potential associates of cognitive performance including (1) self-reported and available modeled dose estimates of environmental exposure, (2) pre-existing and current life stressors, and, in the case of the deployed sample, (3) Gulf experiences.

Preliminary or Pilot Studies.

A. Comparison of Neuropsychological Function in Treatment-seeking and Non-Treatment Seeking PGW Veterans

The objective of this study was to compare the health status, environmental exposure history, and neuropsychological test results of treatment seeking and non-treatment seeking PGW veterans and to obtain preliminary data on the relation of neuropsychological functioning to the presence of health symptoms in the non-treatment seeking group.

A sample of treatment seeking PGW veterans (<u>n</u> = 27) received clinical neuropsychological evaluations after being referred to the neuropsychology service of the Boston DVAMC from clinical services such as neurology, psychiatry, vocational rehabilitation, and general medicine. The most common referral question was "rule out Gulf War Syndrome." The remaining referral questions were, "assess cognition," "rule out memory disorder," and "rule out disorders of attention and concentration." It is likely that all of these questions reflect similar concerns occasioned by patient complaints of CNS symptoms or the physician's impression that the patient might be experiencing the effects of CNS damage. As part of their neuropsychological evaluation patients received a semi-structured clinical interview, including questions concerning health symptoms.

A sample of non-treatment seeking PGW veterans was obtained from among those recruited to date for an ongoing research study from among a cohort of approximately 3,000 military personnel deployed to the PGW from Fort Devens, MA, and followed longitudinally since their return from the Gulf by Dr. Jessica Wolfe and Dr. Susan Proctor. The majority of the cohort had been administered a 20-item Health Symptom Checklist as part of an evaluation 18-24 months after their return. Analysis of these data indicated a median number of participants (51%) reported greater than 5 symptoms. This value was used to stratify the cohort into high symptom (5 or more symptoms) and low symptom (fewer than 5 symptoms) subgroups. A random sample of 24 high and 28 low symptom, non-treatment seeking, PGW veterans was selected for comparison with the clinical group. This sample will be referred to here as the comparison research group.

All participants received an environmental interview, a clinical interview, psychological assessment measures and a brief neuropsychological test battery chosen from a more extensive battery that has been employed by Dr. White in prior work with subjects being assessed for toxicant-induced brain damage from environmental or occupational exposure (Binder, 1993). All of these tests have reliable psychometric properties and have been widely used for both research and clinical purposes. The tests are drawn from two major categories: (1) tests tapping relatively stable, premorbid cognitive/intellectual abilities and (2) tests shown to have high specificity and sensitivity for detecting changes in localized or diffuse cognitive brain functions (e.g., certain forms of attentional skill). Tests in the first category provide uniform estimates of baseline "premorbid" abilities for all individuals; the more specialized tests assess functions that may be differentially affected by a particular disease or condition such as those resulting from exposure to neurotoxicants (White et al., 1994).

The treatment seeking and both high and low symptom comparison research subgroups were comparable in age, education, and gender. On the majority of neuropsychological tests, differences between treatment seeking and both comparison groups were non-significant after adjustment of scores for differences in age, gender, and education. There were, however, several significant differences between treatment seeking veterans and comparison research subjects low in health symptoms. These differences were observed in tests of complex attention (Trails B), motor speed (Finger Tapping with both dominant and non-dominant hands), and learning (California Verbal Learning Test). The treatment-seeking group also reported being more angry, confused, depressed, tense, and tired on the Profile of Mood States (POMS) than the low symptom comparison group. Comparisons between the treatment-seeking group and the high symptom research group yielded fewer differences. The treatment-seeking group made more errors on Trails B and tended to be slower with the dominant hand on a test of manual dexterity (Pegboard). In addition, the treatment-seeking group showed mild retrieval deficits on long-delay free recall on the CVLT compared to the high symptom comparison group. The treatment-seeking group reported being more angry, confused, depressed, tense, and tired on the POMS than the non-treatment seeking high symptom group.

These data suggest that treatment-seeking PGW veterans demonstrate a restricted set of impairments in cognitive abilities compared to non-treatment seeking PGW veterans with few health complaints. Differences in neuropsychological functioning between treatment-seeing PGW veterans and those non-treatment seeking veterans who reported relatively more health complaints were fewer in number. These findings suggest a relation between neuropsychological deficits and the presence of health symptoms in veterans of the PGW. The small number of neuropsychological differences that were observed when treatment-seekers were compared with non-treatment seekers with relatively high numbers of health symptoms may relate to differences in the number or type of symptoms in these groups, or may reflect other differences between treatment-seekers and non-treatment-seekers.

Diagnostic Outcomes in Non-Treatment-Seeking PGW Veterans as Compared With a Non-Gulf
 Deployed Control Group

The objectives of this study conducted by White and colleagues (in press) were to determine the type and frequency of health symptoms in non-treatment seeking PGW veterans, to determine the frequency with which this group met criteria for several neurobehavioral diagnoses, to evaluate their neuropsychological functioning, and to obtain data on their environmental exposure history to relate to these findings.

The study population was selected from a larger cohort of PGW veterans according to an <u>a priori</u> sampling scheme. Sampling was devised to equally represent high and low symptom groups from the Ft.

Devens cohort of PGW-deployed veterans (as previously described) and the group was oversampled for women. Preliminary analyses of 220 PGW deployed and 50 Germany deployed veterans are reviewed below.

Subjects received the same protocol as described in the first study. Subjects included random samples of earlier survey responders from the Devens cohort and a group of Germany deployed veterans (National Guard medical unit from Bangor, ME, deployed to Germany at the time of the PGW). Comparisons of Gulf and Germany deployed groups, adjusted for sampling design, age, education, and gender were made using

SUDAAN. Additionally, comparisons were done of Gulf and Germany deployed groups, controlling for covariate effects such as age, education, gender, race, pre-existing medical concerns, occupational exposures, PTSD or other psychiatric diagnoses, and deployment status.

Preliminary analyses indicated the following health symptoms, in order of frequency: aches and pains, skin rash, fatigue, headache, intestinal problems, and difficulties with memory and concentration. Affective disorders were diagnosed as follows: 7.3% met criteria for current Major Depressive Disorder; 21.6% for lifetime Depressive Disorder, approximately 5% for current PTSD, and 8% for lifetime PTSD.

Analyses of the neuropsychological data indicated differences between PGW deployed veterans and Germany deployed veterans in the areas of motor functioning, mood, and learning. Specifically, differences existed after controlling for covariates including PTSD and Major Depression on measures of simple tracking, memory, and mood. Subtle differences were also found on measures of mood and attention between those Gulf-deployed subjects reporting pesticide exposure and those reporting no exposure, after controlling for covariates, including PTSD and Major Depression.

These studies established the feasibility and reliability of the methodology for the present study. It was found that differences exist in the areas of motor functioning, mood, and learning between Gulf-deployed and non-deployed veterans in a cross-sectional study. In addition, the veterans with neurotoxicant exposures differ from veterans with no exposures on measures of mood and attention.

3. Neuropsychological functioning in treatment-seeking PGW-era veterans

The objective of this study was to compare the health status, environmental exposure history, and neuropsychological test results of treatment seeking PGW-era veterans who were deployed to the Gulf (Research PGW-deployed) and treatment seeking PGW-era veterans who were not deployed to the Gulf (Research PGW-non-deployed). Additional analyses were conducted using a clinically referred PGW-

deployed patient group as a comparison (Clinical-deployed). A sample of treatment seeking PGW-era veterans who were seeking treatment or diagnostic evaluation for any type of health or adjustment complaint were the research subjects for this study. Subjects were chosen because they were listed as having been treated at any of the inpatient or outpatient treatment clinics at the Boston VA, the Manchester VA, the Lowell VA outpatient clinic or the Causeway Street Outpatient Clinic. A group of PGW-deployed veterans who were referred specifically for neuropsychological evaluations because of increased health symptoms were also included. Subjects received the same protocol as described above. Comparisons of Gulf deployed and non-gulf deployed veterans were conducted. Analyses were completed controlling for age and education. Preliminary analyses of the neuropsychological data indicated that differences exist between treatment seeking PGW-deployed veterans and treatment seeking non-deployed veterans when controlling for age and education. Differences exist in the domains of mood, short-term memory, and attention and executive system functions. Specifically, individuals who were deployed to the Gulf reported being more fatigued, had more difficulty acquiring and retaining verbal information, and made significantly more errors on tracking tasks. This study adds further support for the establishment of the feasibility and reliability of the methodology for the proposed study. This study also documents that there are differences in cognitive findings in individuals who were deployed to the Gulf versus their treatment seeking counterparts.

4. A Re-examination of Cognitive Functioning in PGW-Era Veterans

A preliminary study of the re-examination of cognitive deficits in PGW-era veterans included five veterans that had been tested twice by our clinic. A semi-structured questionnaire was administered and the veterans reported a significant decline in terms of health symptoms and cognitive function in the domains of short-term memory and mood. A preliminary analysis of change scores showed variability in performance

from initial testing to follow-up. However, given the limited amount of data, analyses of predictor variables was not completed.

RESEARCH DESIGN AND METHODOLOGY

A. Purpose

The preliminary studies described above indicate that there is a relationship between the health status of PGW veterans and neuropsychological functioning and that these deficits persist over time. What is not known is whether these deficits are progressive. While CNS symptoms are common among PGW veterans' health complaints, the contribution of service in the PGW to CNS symptomatology has remained unclear. The evaluation of CNS dysfunction over time, and, more specifically, the etiology of any progressive cognitive deficits that can be identified, is the primary purpose of the present study. The secondary purpose is to relate the patterns of chronic neuropsychological impairments seen to other variables. We are also interested in understanding the prolonged impact of several variables, including environmental exposures, pre-existing and post-deployment life stressors, psychiatric disorders, existence of multiple health complaints and motivational factors on the study outcomes.

The four preliminary studies described above have indicated that there is a relationship between the health status of PGW veterans and neuropsychological functioning: veterans with greater numbers of health symptoms showed a greater number of neuropsychological impairments than did those with fewer symptoms. This suggests a CNS component to their health complaints. However, these findings leave open the contribution of service in the PGW to these findings. The evaluation of this contribution is the primary purpose of the proposed study. The secondary purpose is to relate the observed patterns of neuropsychological impairments to other variables. The specific objective is to observe and determine whether there are differences in cognitive performance, current health symptom reports, and diagnostic outcomes between the groups of PGW-era veterans. We are also interested in understanding the impact

of several variables, including environmental exposure, pre-existing and post-deployment life stressors, and Gulf experiences on the study outcomes.

B. Study Design

1. Overview

PGW-era veterans who were tested previously in our laboratory are the participants in this study. A group of patients who were initially referred for clinical neuropsychological evaluations (clinical sample) have be re-examined. In addition, a group of individuals who were seeking treatment or diagnostic evaluation for any purpose and who were tested previously (DoD DAMD 17-96-1-6043) are also included in the study sample. As in our pilot studies, test scores from individuals who were deployed to the Gulf are compared with test scores from those who were not deployed. Additionally, longitudinal analyses are being conducted in order to examine changes in test performance. The assessment instruments employed in the study of treatment seekers are those utilized in the preliminary studies described above.

Study Sample

One hundred and fifty subjects will be examined in three years. Subjects are recruited on the basis of the original test date and are tested approximately 4 years (+/- 6 months) after the last neuropsychological evaluation. The recruitment procedure is outlined below. To be consistent with the original recruitment pattern, 75 participants would need to be recruited in the year 1 of the study followed by 50 participants in year 2 and 25 participants in year 3.

Initial Contact. The first contact with potential participants occurs by telephone (or by mail, if there is no telephone number available). The initial contact consists of a description of the follow-up study, reminding subjects of the types of assessment, time required, and reimbursement for their time and effort.

Subjects have an opportunity to ask questions about the procedures. They are also reminded to not inform the

examiners of their Gulf deployment status until the conclusion of the neuropsychological evaluation. They are informed that whether or not they participate will have no bearing on their medical care and that, if they choose to participate, they may withdraw at any time without prejudice. Potential participants are asked to indicate whether they wish to participate, wish not to participate, or wish to defer this decision. In the latter case they are asked whether we may contact them again to determine their decision. Those consenting to participate are asked questions to determine whether they meet preliminary inclusion criteria for the study (that is, that they are not currently in treatment for alcohol or other substance abuse, do not have sensory or motor impairments precluding use of the computer, and did not sustain serious brain injury since the last test session). Prior substance abuse and current medications are recorded but do not constitute exclusion criteria. An appointment at the Boston DVAMC or during one of the field trips is scheduled for patients agreeing to participate.

Veterans retained in the study sample are presented the study consent form for signature.

3. Assessment Instruments and Procedures

a. Behavioral Inventories

1. Cognitive Assessment.

A tester who is blind to the deployment status of the subject administers the neuropsychological test battery. The neuropsychological test battery assesses the functional domains of general intelligence, attention, executive abilities, motor function, visuospatial skills, memory, and mood. The battery is described in detail in Table 1. It includes 1) tests designed to tap relatively stable native intellectual abilities (e.g., WAIS-R Information subtest), and 2) tests shown to have high specificity and sensitivity for detecting changes in neuropsychological functions that have in past studies demonstrated utility in the assessment of toxicant-induced brain damage, and psychiatric disorders (White et al., 1992).

The test battery includes one computer-assisted test from the Neurobehavioral Evaluation System (NES; Letz, 1993), an instrument widely used in the field of occupational health, that represent adaptations of

traditional neuropsychological instruments for computerized stimulus presentation and recording of responses.

The NES instruments have reliable psychometric properties and, as previously described, have demonstrated validity in epidemiological and laboratory studies of exposure to a wide variety of neurotoxicants (Letz, 1993).

Because it is anticipated that issues related to compensation for disability may be pertinent for many of these veterans who may also be treatment seeking and compensation seeking, the possibility of motivational factors affecting test performance (Binder, 1992; Binder, 1993) are addressed directly. This is accomplished in two ways: 1) by inclusion of internal consistency checks in certain neuropsychological test measures (e.g., test/item repetition, easy/hard item comparisons) and evaluation of response style (over-endorsement, response approximations) and 2) by administering the Test of Motivation and Malingering (TOMM; Tombaugh, 1996).

Table 2. Full Neuropsychological Test Battery; Tests in Preliminary Studies Marked *

I. Tests of General Intelligence

Test Name

Description

1. Wechsler Adult Intelligence Scale-Revised Information usually learned in school; (WAIS-R)(53)

to assess native intellectual abilities

Information subtest

II. Tests of Attention, Vigilance, and Tracking

Test Name	Description		
1. Digit Span Subtests: WAIS-R*	Oral recall of digits forward and backward; Given twice to assess		
Wechsler Memory-Scale-Revised (WMS-R)	consistency of performance		
2. Trail Making Test*	Timed connect-a-dot task to assess attention and motor control requiring sequencing (A) and alternating sequences (B)		
2. Continuous Performance Test (CPT)*	Target letter embedded in series of distracters; to assess sustained attention and reaction time		
 Paced Auditory Serial Addition Test * (PASAT;) 	Continuous addition test; to assess sustained attention under distracting conditions		
5. Wisconsin Card Sorting Test* (WCST)	Requires use of feedback to infer decision making rules; assesses problem solving ability and flexibility		

III. Tests of Motor Function

cription

1. Finger Tapping*

Speed of tapping with index finger of each hand assesses simple motor speed

2. Purdue Pegboard Test*

Speed of inserting pegs into slots using each hand separately and both together

IV. Tests of Visuospatial Function

Test name

Description

1. WAIS-R Block Design*

Using blocks to replicate 2-dimensional designs; to assess visuoconstruction

2. Hooper Visual Organization Test *

Identifying objects from line

(HVOT)

drawings of disassembled parts; assesses ability Synthesize visual stimuli

V. Tests of Memory

Test Name

Description

1. Paired Associate Learning-WMS-R* Recall of second member of pairs of words immediately and after a delay; assesses short-term memory and recall

2. Visual Reproductions-WMS*

Reproductions of visual designs, immediately and after a delay; assesses short term memory and recall

3. California Verbal Learning Test*

(CVLT)

List of 16 nouns from 4 categories presented over multiple learning trials with recall after interference

VI. Tests of Personality and Mood

Test Name

Description

1. Profile of Mood States* (POMS)

65 single word descriptors of affective symptoms are endorsed for degree of severity and summed on six mood scales

VII Tests of Malingering

Test Name

Description

1. Test of Motivation and Malingering* (TOMM)

Immediate forced choice recognition of line drawings of 50 common objects; assesses motivation and malingering

2. Psychological Assessment.

- 1) Participants are administered the Structured Clinical Interview for DSM-III-R (SCID, Spitzer et al, 1990) and a current Global Assessment of Functioning is assessed. This instrument has demonstrated reliable psychometric properties for determining the presence or absence of current or past major Axis I disorders. Dr. Krengel also re-administers the Clinician Administered PTSD Scale (CAPS; Blake, 1990a), a state-of-the-art instrument for confirming the diagnosis of current or past PTSD and for evaluating the intensity, frequency, and severity of the disorder and its individual symptom criteria. Extensive research now indicates that this instrument has highly acceptable psychometric properties (Weathers et al., submitted). Dr. Krengel is not aware of the subject's prior responses on either measure before conducting the interviews. Study participants fill out a series of self-report, paper and pencil measures designed to confirm and define symptoms of PTSD (Mississippi Scale for PTSD; Keane 1988), to describe war-zone exposure (modified Laufer Combat Scale; Gallops, 1981), and to identify traumatic events occurring since last assessment.
- 2) Dr. Krengel also conducts a semi-structured clinical interview eliciting information pertaining to recent past and current mood disorders, substance use, neurological and medical illness, traumatic brain injury, and history of other traumatic events. Study participants are also asked questions specifically related to recent occupational history (including possible occupational exposure to neurotoxicants), family history of psychiatric disorder, and life stressors since the time of the last evaluation. Any treatment programs initiated since the last assessment are documented.
- 3) Study participants complete a symptom checklist consisting of a comprehensive list of frequently reported health and mental health symptoms, which is compared to their previous questionnaire data.

b. Environmental Exposure Assessment

For the purpose of this study, the environmental interview used in previous studies (Wolfe et al., 1998) has been modified and put in questionnaire form. The questionnaire includes questions pertaining to recent occupational and environmental exposures. The original semi-structured interview schedule is modified from

those in current use and includes details of past military and health experiences, pre-military environmental exposure and health. Included is length of stay, geographical location, unit/individual duties, movement in the theater, and environmental exposure during deployment (type, intensity, frequency, duration, locale). The environmental health specialist (Dr. Susan Proctor) is available to interview subjects in the event that questionnaire data concerning environmental and occupational exposures need clarification or when a more extensive interview is required.

This study also makes use of objective data on geographical locations and cumulative exposure <u>at the unit level</u> for the clinical subjects (obtained from D. Hakenson, Center for Research on Unit Records, [RUR], U.S. Army, personal communication). RUR is directly involved with the determination and computerization of all U.S. military locations during the war, including grid coordinates and longitude/latitude demarcations. This information is linked with existing environmental data (e.g., concentrations of oil smoke particulates modeled by the U.S. Army Center for Health Promotion and Preventative Medicine (CHPPM)). With input from this team, we will seek information on objective exposure measures and will integrate these data with the geographical locations reported by our study participants to better estimate individual cumulative exposure for the sample of participants deployed to the Gulf.

Data Analysis

Treatment of Data

Preliminary analyses will be conducted to check for baseline differences in those who have agreed to participate and those lost to follow-up. A comparison of treatment seekers versus the clinical group on education, combat exposure, and other characteristics of interest will be completed. In addition, we will explore the relationship of symptom reporting and diagnostic outcome between the groups and its correlation with change in neuropsychological test performance.

The aims of this study are to determine whether there has been a diminution in performance on neuropsychological tests from the initial testing to follow-up in the PGW sample and to explore subject characteristics and individual risk factors in Gulf exposure in accounting for this change.

- a. Analyses will focus on **difference** scores describing change in performance in follow-up and initial test performances. A crude comparison between the average change in performance for the Gulf-deployed versus non-deployed group will be conducted through a paired sample <u>t</u>-test. A comparison of PGW-deployed and non-deployed groups on average change in performance adjusting for intervening factors will be conducted through analysis of covariance.
- b. In addition, we will split the PGW sample into treatment seekers and the clinical sample and make a comparison across the 3 groups using a 1-factor analysis of variance on the average difference scores. We will examine baseline characteristics that might be related to change in neuropsychological performance and examine change in intervening events or stressors that might relate to change in performance.

For another aim of this study, focusing on the role of neurotoxicant exposure in the Gulf, we will examine the relationship between experience in the Gulf and change in neuropsychological performance through multiple regression. These analyses will include indicator variables to account for group status (1= Gulf deployed clinical sample, 2= Gulf deployed treatment seekers, 3= non-deployed) as well as individual risk factors and intervening risk factors that might be related to outcomes. Additional analyses exploring interactions between exposures and change in performance will be pursued. We will look at the relationship of stress, psychiatric diagnoses and disability status, and normal aging through multiple regression analyses as described above.

KEY RESEARCH ACCOMPLISHMENTS

STATUS/RESULTS TO DATE

The objective of this study is to determine whether veterans of the Gulf War (GW) show cognitive impairments suggestive of central nervous system damage at a greater rate than do GW-era veterans who were not deployed to the Gulf. A further study objective is to assess whether the GW veterans' symptoms are actually becoming worse since their first evaluation approximately four years ago. The diagnostic protocol for these evaluations includes an environmental exposure questionnaire, a psychological assessment and a neuropsychological evaluation. The environmental questionnaire is overseen by an environmental health specialist (Dr. Proctor) and assesses health experiences, geographical location, unit/individual duties, and environmental exposures during deployment to the Gulf (type, intensity, frequency, duration, locale), and/or during military service in general. In addition, subjects are queried about current health symptomatology including evaluation of chronic fatigue syndrome and multiple chemical sensitivities. The psychological assessment is administered by a clinical psychologist (Dr. Krengel) and includes the administration of the Structured Clinical Interview for DSM-IV (SCID; Spitzer et al., 1990), the Clinician Administered PTSD scale (CAPS-DX; Blake et al, 1995) and a semi-structured clinical interview. The SCID interview is reliably used to determine Axis I disorders (Wolfe & Keane, 1993) and the CAPS-DX is a state of the art instrument for confirming the diagnosis of current and/or lifetime PTSD, as well as, for evaluating the intensity, frequency and severity of the disorder and its individual symptom criteria. The semi-structured clinical interview administered by Dr. Krengel elicits information pertaining to current or past mood disorders, substance abuse, neurological and medical illness, traumatic brain injury, history of other traumatic events, birth trauma, developmental delays, and history of learning disability. This interview includes a follow-up of any medical or social changes since the study participant's last evaluation. Subjects are asked questions specifically related to changes in school achievement, and occupational history (including possible occupational exposure to neurotoxicants), family history of psychiatric disorder, and post-deployment

stressors. The neuropsychological test battery is administered by trained professionals. The total time to complete the study protocol is approximately 3-4 hours.

This longitudinal study is still in the data collection phase and our enrollment at this time includes 44 study participants from the original randomized list of Persian Gulf War-era veterans. Of this group, 41 subjects are GW veterans and 3 subjects are non-PGW veterans. Specifically, of the 41 GW veterans recruited, 11 were treatment-seekers and 30 were from the clinical sample. Our initial projections for year 1 recruitment included 75 study participants. The focus of recruitment for year 1 was GW veterans and not control veterans because the majority of control veterans were not recruited in the first years of the original study and therefore, have not met the predetermined time period (4 years) between evaluations. In terms of recruitment for GW veterans, a slow start-up phase and minor nonlocation difficulties in recruiting brought our recruitment total for year1 below our initial projections. Although our total participant recruitment for year 1 was below our initial projections, those participants that we have been able to reach by telephone have been largely receptive to our study and our dropout rate is below ten percent. We have employed several new strategies to improve our recruitment efforts including updating residence locations through a federal interagency agreement, performing national yellow page searches on the internet and contacting reported next of kin. These strategies have improved our study participant contact ratio and will allow us to improve our recruitment efforts in order to meet our goal of 150 study participants.

In terms of reportable outcomes, some preliminary results have been submitted for two poster presentations at the American Psychological Association annual meeting in San Francisco, CA in August, 2001. We performed some preliminary analyses after obtaining data from our first 24 study participants. The analyses involved paired sample t-tests comparing participants' performances at Time 1 and Time 2 evaluations. These preliminary analyses suggested relatively static test-retest performances in we weterans within a four-year time interval. These preliminary findings do not support pentinued

diminution of veteran's performances on neuropsychological measures. However, neither do these performances suggest improvement of cognitive functioning over this time period. Preliminary analyses of mood changes in the first 24 study participants were also conducted using paired sample t-tests and suggested that current mood-state has changed during the past four years. Specifically, mood complaints were less robust with veterans reporting lower levels of tension, anger and confusion. However, the GW veterans reported equivalent levels of fatigue when compared with their original assessments.

FUTURE DIRECTIONS

This sample was also examined using the same measures as a sample already studied by the investigators (BEHC Ft. Devens sample, n = 220), who were seen as part of a cohort study of deployed PGW veterans. Using data from these two samples, it is now possible to examine separately the effects of treatment seeking, deployment, diagnosis, and PGW exposures on various outcome measures.

When comparing the original testing of the PGW-deployed treatment seeking sample (DoD) to the largely non-treatment seeking sample (BEHC), it was found that treatment seeking veterans performed worse on a significant number of cognitive tasks (in the functional domains of attention, executive skills, fine manual motor dexterity, visual constructive ability, and short-term memory). They were also more dysphoric on mood scales. The current study will help assess if these patterns continue in the treatment-seeking cohort four-years after their first evaluations.

In our original evaluations of treatment seekers, deployment status (deployed versus non-deployed) had no effect on likelihood of being diagnosed with PTSD (13.2% deployed, 10.8% non-deployed), or MCS or CFS (<1% in all groups). The current follow-up study will continue to assess for these syndromes and report any changes in diagnostic incidence.

In our original evaluations, PTSD diagnoses and deployment status were associated with poorer performance on neuropsychological tests in three domains (attention, visual construction, memory) and with greater mood complaints, though the findings were less pronounced than those seen when comparing treatment

seekers to non-treatment seekers. These findings raise the possibility that treatment seekers are experiencing CNS dysfunction from some cause that is not unknown. PTSD is one candidate, but this diagnosis does not explain all of the findings. In the DoD (treatment-seeking) deployed sample, self-reported exposure to pesticides, burning human waste with diesel fuel and PB were considered as explanations of CNS dysfunction, but exposure-outcome relationships were not confirmed. We will continue to assess for these relationships in the current follow-up examination. However, due to the small incidence of these diagnoses and potential for Type II error, it seems appropriate to perform these analyses when more study participants have been recruited.

Reportable outcomes

Abstracts have been accepted for presentations at scientific meetings. In addition, one manuscript has been accepted pending revisions.

Abstracts:

- Neuropsychological findings among Persian Gulf War-era veterans, presented at International Neuropsychological Society meeting in Honolulu, Hawaii, February 1998.
- Relationship of PTSD and depression to Persian Gulf health problems, presented at the Conference on Federally Sponsored Gulf War Veterans Illnesses research meeting in Washington, D.C., June 1998.
- Neuropsychological Findings Among Persian Gulf War-era veterans, presented at the Conference on Federally Sponsored Gulf War Veterans Illnesses research meeting in Washington, D.C., June 1998.
- 4. Neuropsychological test methods in assessment of neurotoxicant exposure in Persian Gulf War-era veterans, presented at the Theoretical and Experimental Neuropsychology annual meeting, June 1998.
- 5. Neuropsychological deficits in treatment-seeking Persian Gulf War-era veterans, presented at the Conference on Federally Sponsored Gulf War Veterans' Illness Research, June, 1999.

- 6. Neuropsychological deficits in treatment-seeking Persian Gulf War-era veterans, presented at the International Neuropsychological Society annual meeting in Denver, Colorado, February, 2000.
- 7. Neuropsychological deficits in association with stress reaction and pyridostigmine bromide intake in Persian Gulf War-era veterans, presented at the International Neuropsychological Society annual meeting in Denver, Colorado, February, 2000.
- 8. Neuropsychological deficits in association with stress reaction and pyridostigmine bromide intake in Persian Gulf War-era veterans, presented at the Conference on Federally Sponsored Gulf War Veteran's Illness Research, January 2001.
- 9. A re-examination of neuropsychological functioning in Gulf War-era veterans, submitted for presentation to the American Psychological Association annual meeting in San Francisco, CA, August, 2001.
- 10. A re-assessment of current mood state in Persian Gulf War-era veterans, submitted for presentation to the American Psychological Association annual meeting in San Francisco, CA, August, 2001.

Paper accepted with revisions: White et al., manuscript submitted entitled, "Neuropsychological function in Gulf War Veterans: Relationships to self-reported toxicant exposures." American Journal of Epidemiology.

Planned manuscripts: Proctor et al., manuscript planned entitled "Comparison of functional status (as measured by SF36) in clinical and research studies of GW-deployed military personnel."

Proctor et al., manuscript planned entitled "Methodology to assess environmental exposures in GW-deployed military personnel."

Krengel et al., manuscript planned entitled "Neuropsychological functioning in PGW-era veterans."

Sullivan et al., manuscript planned entitled "Neuropsychological deficits in association with stress reaction and pyridostigmine bromide intake in PGW-era veterans."

Conclusions:

The preliminary analyses of the test-retest measures for the first 24 study participants suggest relatively static cognitive performances and slightly less robust mood complaints in GW veterans within a four-year time interval. These preliminary findings do not support continued diminution of veteran's performances on neuropsychological measures. However, neither do these performances suggest improvement of cognitive functioning over this time period. It is possible that this relatively static performance in our GW veterans could be consistent with long-term patterns expected from neurotoxicant exposure. However, continued follow-up of cognition and mood and a full study sample should provide a clearer picture of this pattern.

The study has the following clinical implications:

- 1). CNS complaints of PGW-era veterans should be taken seriously and explored with formal testing and neuropsychological diagnosis.
- 2). Mood complaints are common among deployed and non-deployed PGW-era veterans.
- 3). PTSD occurs in PGW-era veterans (10-13%) and may be related or unrelated to the war. Possibility of its existence should be considered.
- 4). PGW exposure to environmental agents should be considered in evaluation of PGW veterans' health.
- 5). Evaluation of CFS and MCS should be done carefully, with established diagnostic criteria.
 CFS- and MCS-like symptoms are probably more common than the full blown syndromes.

REFERENCES

- Almog, S., Winkler, E., Amitai, Y., Dani, S., Shefi, M., Tirosh, M., & Shemer, J. (1991). Acute pyridostigmine overdose: A report of nine cases. Israel Journal of Medical Science, 27, 659-663.
- Andrews, L. S. & Snyder, R. (1986). Toxic effects of solvents and vapors. In C. D. Klaassen, M. W. Amdur, & J. Doull (Eds.), <u>Casarett and Doull's toxicology: The basic science of poisons</u> (3rd ed.). New York: Macmillan.
- American Psychiatric Association. (1995). <u>Diagnostic and statistical manual of mental disorders</u>, <u>DSM-IV</u>. Washington, DC: Author.
- Anger, W.K. (1990). Worksite behavioral research. Results, sensitive methods, test batteries and the transition from laboratory data to human health. <u>Neurological Toxicology</u>, <u>11</u>, 629-720.
- Axelrod, B.N., and Milner, I.B. (1997). Neuropsychological Findings in a Sample of Operation Desert Storm Veterans. <u>Journal of Neuropsychiatry and Clinical Neuroscience</u>, 9:23-28.
- Baker, E.L., Feldman, R.G., White, R.F., Harley, J.P., Niles, C.A., Dinse, G.E., Berkey, C.S. (1984).

 Occupational lead neurotoxicity: A behavioral and electrophysiological evaluation. Study design and year one results. <u>British Journal of Internal Medicine</u>, <u>41</u>, 353-361.
- Binder, L.M. (1992). Forced-choice testing provides evidence of malingering. <u>Archives of Physical and Medical Rehabilitation</u>, 73, 377-380.
- Binder, L.M. (1993). Assessment of malingering after mild head trauma with the Portland Digit Recognition Test. <u>Journal of Clinical and Experimental Neuropsychology</u>, <u>15</u>, 170-182.
- Black, P.H. (1994). Central nervous system-immune system interactions: psychoneuroendocrinology of stress and its immune consequences. <u>Antimicrob-agents-chemother</u>. 38 (1) 1-6.
- Blake, D., Nagy, L., Kaloupek, D., Klauminzer, G., Charney, D., & Keane, T. (1990a). A clinician rating scale for assessing current and lifetime PTSD: The CAPS-1. The Behavioral Therapist 18, 187-188.

- Blake, D., Weathers, F., Nagy, L., Kaloupeck, D., Klauminzer, G., Charney, D., & Keane, T. (1990b).
 <u>Clinician-administered PTSD scale for DSM-IV (CAPS-DX)</u>. National Center for Posttraumatic Stress
 Disorder, Behavioral Science Division, Boston and Neurosciences Division, West Haven.
- Burchfiel, J.L. & Duffy, F.H. (1982). Organophosphate neurotoxicity: Chronic effects of sarin on the electroencephalogram of monkey and man. <u>Neurobehavioral Toxicology & Teratology</u>, <u>4</u>, 767-778.
- Chevillard, M., Lainee, P., Robineau, P., & Puchelle, E. (1992). Toxic effects of sulfur mustard on respiratory epithelial cells in culture. <u>Cell Biology and Toxicology</u>, <u>8</u>, 171-181.
- Cullen, M.R. (1987). The worker with multiple chemical sensitivities: an overview. In M. Cullen (Ed.), Workers with Multiple Chemical Sensitivities (pp. 655-662). Philadelphia, PA: Hanley & Belfus.
- Doucet I. (1994). Desert storm syndrome: sick soldiers and dead children? Medicine and War 10:183-194.
- Echeverria, D. & White, R F. (1992, September). A neurobehavioral evaluation of PCE exposure in patients and dry cleaners: A possible relationship between clinical and pre-clinical effects. Paper presented at the Ninth International Symposium of Epidemiology in Occupational Health, Cincinnati, OH.
- Eisen, E.A., Letz, R, Wegman, D.H., Baker, E.L., & Pothier, L. (1988). Defining measurement precision for effort-dependent neurobehavioral tests: The case of three neurobehavioral tests. <u>International</u>

 <u>Journal of Epidemiology</u>, 17, 292-298.
- Everson, M.P., Kotler, S., Blackburn, W.D. (1999). Stress and immune system dysfunction in Gulf War veterans. Annals of the New York Academy of Sciences, 876, 413-418.
- Friedman, A., Kaufer, D., Shemer, J., et al. (1996). Pyridostigmine brain penetration under stress enhances neuronal excitability and induces early immediate transcriptional response. <u>Nature Medicine 2(12):1382-1385</u>.
- Fukuda, K., Straus, S.E., Hickie, I., et al. (1994). The chronic fatigue syndrome: A comprehensive approach to its definition and study. <u>Annals of Internal Medicine</u>, <u>121</u>, 953-959.

- Gallops, M., Laufer, R.S., & Yager, T. (1981). The combat scale: Revised. In A. Egendorf, C. Kadushin, R.S. Laufer, G. Rothbort, & L. Floan (Eds.), <u>Legacies of Vietnam: Comparative adjustment of veterans and their peers</u>. New York: Center for Policy Research.
- Goldstein, G., Beers, S.R., Morrow, L.A., et al. (1996). A preliminary neuropsychological study of Persian Gulf veterans. <u>Journal of the International Neuropsychological Society</u>, 2:368-371.
- Gray, G.C., Coate, B.D., Anderson, C.M. et al. (1996). The Postwar hospitalization experience of U.S. veterans of the Persian Gulf War. New England Journal of Medicine. 335:1505-13.
- Haley, R.W., Kurt, T.L., and Horn, J. Is there a Gulf War Syndrome?: Searching for syndromes by factor analysis of symptoms. <u>Journal of the American Medical</u> <u>Association</u> (1997) 277 (3):223-230.
- Haley, R.W. Is Gulf War Syndrome due to stress? The evidence reexamined. (1997). <u>American Journal of Epidemiology</u>. 146(9): 695-703.
- Hanninen, H., Hernberg, S., Mantere, P., Vesaito, R. & Jalkaner, M. (1978). Psychological performance of subjects with low exposure to lead. <u>Scandinavian Journal of Work and Environmental Health</u>, 2, 240-255.
- Hom, J., Haley, R.W., Kurt, T.L. (1997). Neuropsychological correlates of Gulf War Syndrome.

 Archives of Clinical Neuropsychology, 12 (6), 531-544.
- Husain, K., Vijayaraghavan, R., Pant, S.C., Raza, S. K., Pandey, K.S. (1993). Delayed neurotoxic effect of sarin in mice after repeated inhalation exposure. <u>Journal of Applied Toxicology</u>, <u>13</u>, 143-145.
- Institute of Medicine (IOM). (1996). Health consequences of service during the Persian Gulf War: Initial findings and recommendations for immediate action. Washington, DC: National Academy Press.
- Joseph, S.C.(1997). A Comprehensive clinical evaluation of 20,000 Persian Gulf War veterans. Military

 Medicine. 162(3):149-155.
- Kadar, T., Shapira, S., Cohen, G., Sahar, R., Alkalay, D., & Raveh, L. (1995). Sarin- induced neuropathology in rats. <u>Human and Experimental Toxicology</u>, 14, 252-259.

- Kawabuchi, M., Cintra, W.M., Deshpande, S.S., & Dlbuquerque, E.X. (1991). Morphological and electrophysiological study of distal motor nerve fiber degeneration and sprouting after irreversible cholinesterase inhibition. Synapse, 8, 218-228.
- Keane, T.M., Caddell, J.M., & Taylor, K.L. (1988). Mississippi scale for combat related post-traumatic stress disorder: Three studies in reliability and validity. <u>Journal of Consulting and Clinical Psychology</u>, 56, 85-90.
- Kellner, M., Baker, D.G., Yehuda, R. (1997). Salivary cortisol and PTSD symptoms in Persian Gulf War combatants.
- Kimerling, R. & Calhoun, K.S. (1994). Somatic symptoms, social support, and treatment seeking among sexual assault victims. Journal of Consulting and Clinical Psychology, 62, 333-340.
- Klaassen, C.D., Amdur, M.O., & Doull, J. (Eds.), (1986). <u>Casarett and Doull's toxicology: The basic</u> science of poisons (3rd ed.). New York: Macmillan.
- Kreutzer, R.D., Grogl, M., Neva, F. et al. (1993). Identification and genetic comparison of leishmanial parasites causing viscerotropic and cutaneous disease in soldiers returning from Operation desert storm. American Journal of Tropical Medicine and Hygiene 49 (3):357-363.
- Kulka, R.A., Schlenger, W.E., Fairbank, J.A., Hough, R.L., Jordan, B.K., Marmar, C.R., & Weiss, D.S. (1990). Trauma and the Vietnam War generation. New York: Brunner/Mazel.
- Lallement, G., Foquin, A., Baubichon, D., Burckhart, M., Carpentier, P., Canini, F. (1998). Heat stress even extreme, does not induce penetration of pyridostigmine into the brain of guinea pigs.

 Neurtoxicology, 19 (6), 759-766.
- . Letz, R., & Baker, E.L. (1988). <u>Neurobehavioral Evaluation System: NES User's Manual.</u> Winchester, MA: Neurobehavioral Systems, Inc.
- Levy, C. (1988). Agent orange exposure and posttraumatic stress disorder. <u>Journal of Nervous and Mental Disease</u>, <u>176</u>, 242-245.
- Lezak, M.D. (1995). Neuropsychological assessment (3rd edition). New York: Oxford University Press.

- Litz, B.T., Keane, T.M., Fisher, L., Marx, B., & Monaco, V. (1992). Physical health complaints in combat-related post-traumatic stress disorder: A preliminary report. <u>Journal of Traumatic Stress</u>, <u>5</u>, 131-141.
- Nicolson, G.L., and Nicolson, N.L. (1996). Diagnosis and Treatment of Mycoplasmal Infection in Persian Gulf War Illness-CFIDS Patients. <u>International Journal of Occupational Medicine</u>, Immunology, and Toxicology 5(1):69-78.
- Oster, C.N., and Sanford, J.P. Febrile illness in a Desert Storm veteran. Presented at conference, "Hospital Practice.", 1992
- Rockefeller, J.D., Deconcini, D., Mitchell, G., Graham, B., et al. Is Military Research Hazardous to

 Veteran's Health? Lessons Spanning Half a Century, <u>United States Senate Committee on Veteran's</u>

 <u>Affairs staff report</u>, 1994.
- Riegle, Senator D.W. (1993, September 9). Gulf war syndrome: The case for multiple origin mixed chemical/biotoxin warfare related disorders. Boston Globe.
- Roy, M.J. Evaluating the symptoms of Persian Gulf War veterans. Presented at conference, "Federal Practitioner." 1994
- Shalev, A., Bleich, A., & Ursano, R.J. (1990). Post-traumatic stress disorder: Somatic comorbidity and effort tolerance. Psychosomatics, 31, 197-203.
- Sillanpaa, M.C., Agar, L.M., Milner, I.B., Podany, E.C., Axelrod, B.N. & Brown, C.G. (1997).
- Gulf War veterans: a neuropsychological examination. J Clin Exp Neuropsychol., 19 (2): 211-19.
- Spitzer, R.L., Williams, J.B. W., Gibbon, M., & First, M.B. (1990). Structured clinical interview for DSM-III-R-non-patient edition (SCID-NP, version 1.0). Washington, DC: American Psychiatric Press.
- Sullivan, K., Krengel, M., White, R.F., S.P. Proctor, Devine, S.P., & Davis, L. (2001). The effects of pyridostigmine bromide and stress on cognitive functioning in Persian Gulf War-era veterans.

 (Abstract), Conference on illnesses among Gulf War Veterans: A decade of scientific research

- Sutker, P.B., Vasterling, J.J., Brailey, K., & Allain, A.N. (1995). Memory, attention, and executive deficits in POW survivors: Contributing biological and psychological factors. Neuropsychology, 9, 118-125.
- Tombaugh, T. (1996). Test of memory malingering. New York: Multi-health systems.
- U.S. Congress, Office of Technology Assessment. (1993). The Department of Veterans Affairs Persian Gulf Veterans' Health Registry. Washington, D: .OTA- H-577.
- Vasterling, J.J., Brailey, K., Constans, J.I. & Sutker, P.B. (1998) Attention and memory dysfunction in Post-traumatic stress disorder. <u>Neuropsychology</u>. 12(1): 125-133.
- Weathers, F.W., Blake D.D., Krinsley, K.E., Haddad, W.H., Huska, J.A., & Keane, T.M. Reliability and validity of the clinician-administered PTSD scale. Manuscript submitted for publication.
- White, R.F., Feldman, R.G., & Proctor, S.P. (1992). Neurobehavioral effects of toxic exposures. In R. F. White (Ed.), Clinical Syndromes in Adult Neuropsychology: The Practitioner's Handbook (pp. 1-51). Amsterdam: Elsevier.
- White, R.F. & Proctor, S.P. (1992). Research and clinical criteria for the development of neurobehavioral test batteries. <u>Journal of Occupational Medicine</u>, <u>14</u>, 140-148.
- White, R. F., Robins, T.G., Proctor, S.P., Echeverria, D., & Rocskay, A.S. (1994). Neuropsychological effects of naphtha exposure among automotive workers. <u>Occupational and Environmental Medicine</u>, 51, 102-112.
- White, R.F., Proctor, S.P., Heeren, T., Wolfe, J., Krengel, M., Vasterling, J., Lindem, K., Heaton, K., Sutker, P., & Ozonoff, D., (submitted) Neuropsychological function in Gulf War veterans: relationships to self-reported toxicant exposures. <u>American Journal of Epidemiology</u>. (Accepted pending revisions).
- Wintermeyer, S.F., Pina, J.S., Cremins, J.E. et al. (1994). The inpatient experience of a U.S. Army combat support hospital in the Persian Gulf during non-combat and combat periods. <u>Military</u>

 <u>Medicine</u> 159:746-751.

- Wolfe, J. & Charney, D.S. (1991). Use of neuropsychological assessment in posttraumatic stress disorder. Psychological Assessment: A Journal of Consulting and Clinical Psychology, 3, 573-580.
- Wolfe, J., Kelley, J.M., Buscela, M.L., & Mark, W.R. (1992). Ft. Devens Reunion Survey: Report of Phase I. In R. Rosenheck et al. (Eds.), <u>War-zone stress among returning Persian Gulf troops: Final Report</u>. Legislative report to Congress.
- Wolfe, J., Brown, P.J., & Kelley, J.M. (1993). Reformulating war stress: Exposure and the Persian Gulf War. Journal of Social Issues, 49, 15-31.
- Wolfe, J. & Keane, T.M. (1993). New perspectives in the assessment and diagnosis of combat-related posttraumatic stress disorder. In J.P. Wilson & B. Raphael (Eds.), International handbook of traumatic stress syndromes. New York: Plenum Press.
- Wolfe, J., Keane, T.M., Gerardi, R.A., Butters, M., Mora, C.A., & Weathers, F. <u>Information processing</u>

 <u>and memory changes associated with combat-related post-traumatic stress disorder</u>. Manuscript submitted for publication.
- Wolfe, J., Proctor, S.P., Davis, J.D., Borgos, M.& Freidman, M.J. (1998). Health symptoms reported by Persian Gulf War veterans two years after return. <u>AJIM</u>, 33, 104-113.
- Yokoyama, K., Araki, S. & Aono, H. (1988). Reversibility of psychological performance in subclinical lead absorption. <u>Neurological Toxicology</u>, 9, 405-410.

Submitted for presentation at the American Psychological Association

Annual meeting at San Francisco, CA August, 2001

A Re-examination of Neuropsychological Functioning in Gulf War-Era Veterans

Introduction:

Existing data from the Boston Environmental Hazard's Research Center suggest that Persian Gulf War (PGW) era veterans deployed to the Gulf have deficits on a limited, specific group of neuropsychological test measures as compared to PGW-era veterans who were not deployed to the Gulf. These findings raise the possibility of subtle, subclinical central nervous system damage associated with Gulf deployment and raise the question of whether such changes may be attributable to exposure to neurotoxicants in the Gulf environment. This latter possibility is particularly an issue because the existence of relative cognitive deficits was related in our sample to self-reported environmental exposures that are likely neurotoxic.

Recently some PGW veterans have reported that their cognitive functioning and mood complaints are worsening over the time since their return from the Gulf. Veterans are specifically reporting further deficits in the areas of short-term memory and attention. For instance, veterans report that they are having more difficulty remembering names and appointments and they are having difficulty completing work-related tasks. These reports are at odds with the usual course of central nervous system (CNS) effects of intoxication, which generally remit or remain static in the absence of exposure.

The specific aim of the project was to determine whether objective neuropsychological test measures reveal any progressive diminution in cognitive function among Gulf deployed PGW veterans by comparing test performance observed initially (1995-1998) with performance 4 years (+/- 6 months) later.

Subjects:

In our initial study, 207 PGW-deployed treatment seeking veterans were administered a thorough neuropsychological test battery including assessment measures from the domains of memory, attention and executive systems, visuospatial, language, motor and mood. Overall, neuropsychological deficits were found in the areas of attention and short-term memory when controlling for age, education and mood. Twenty-four individuals from our original GW deployed sample have been tested for a second time approximately four years after their first evaluation.

Preliminary Results:

Results of paired T-tests indicated similar performances on tests from most domains including attention and executive functions. The only significant finding was in the domain of nonverbal memory, specifically the WMS-R visual reproductions delayed recall. This performance improved significantly relative to the first testing.

Summary:

These preliminary results from our first 24 subjects suggest relatively static performances in GW veterans on test-retest measures with a four-year interval. These preliminary findings do not support continued diminution of veteran's performances on neuropsychological measures. However, neither do these performances suggest improvement of cognitive functioning over this time period. These results will be discussed as they relate to etiologic factors including possible neurotoxicant exposures. It is possible that this relatively static performance in our GW veterans could be consistent with long-term patterns expected from neurotoxicant exposure. However, continued follow-up of cognition and mood in these veterans should provide a clearer picture of this pattern.

Submitted for presentation at the American Psychological Association annual

meeting in August, 2001, San Francisco, CA

A Re-assessment of current mood state in Persian Gulf War-Era Veterans

Introduction:

It has been found that Persian Gulf War (PGW) deployed veterans have reported a variety of mood complaints including depression, tension, anger and confusion since their return from the Gulf. These mood complaints have been reported to effect performance at work and have impacted social functioning, according to many PGW-deployed veterans. Some researchers have also found that these mood symptoms have had an effect on performance on a limited number of neuropsychological tests. Existing data from the Boston Environmental Hazard's Research Center suggests that PGW veterans deployed to the Gulf have significantly more mood complaints than their non-deployed counterparts. Although these complaints coexist with impairments in specific cognitive domains, we have found that neuropsychological deficits persist when controlling for current mood state. Given the impact of current mood on daily functioning, it is important to monitor mood state in PGW-deployed veterans in order to assist individuals in developing strategies to enhance social and occupational functioning.

We are currently following a clinical sample of over 200 PGW-era veterans, over half of whom were deployed to the Gulf during the war. Clinically, many of the PGW- era deployed veterans continue to report mood changes that have reportedly magnified over time since their return from the Gulf. Specifically, PGW-era deployed veterans report increased amounts of tension, depression and fatigue during clinical interview at a significantly higher rate than their non-deployed peers.

The specific aim of the current project was to determine whether mood measures revealed any progression of mood complaints among Gulf deployed PGW veterans by comparing scores observed initially (1995-1998) and those 4 years (+/- 6 months) later.

Subjects:

In our initial study, 207 PGW-deployed treatment seeking veterans were administered a thorough neuropsychological test battery including assessment measures from the domains of memory, attention and executive systems, visuospatial skills, language, motor speed and dexterity, and mood. PGW-era deployed veterans complained of fatigue, tension, depression, anger and confusion when compared to PGW-era nondeployed veterans. Twenty-four individuals from our original PGW-era deployed sample have returned to our clinic for follow-up. These individuals were interviewed and assessed for a second time approximately four years after their first evaluation.

Preliminary Results:

Results of paired T-tests indicated equal results for fatigue over four years in the deployed group, whereas levels of tension and anger diminished slightly relative to initial assessment. Confusion declined significantly from prior reports (p = .05). The only clinically significant mood complaint in this group of individuals is fatigue (T = 58).

Summary:

These preliminary results from our first 24 subjects suggest that current mood state has changed over time in PGW veterans on test-retest measures over a four-year interval. Specifically, mood complaints have become slightly less robust. These results will be discussed as they relate to follow-up neuropsychological test performance and social and occupational functioning in this group of PGW-deployed veterans.